

Colonoscopy Prep using Clenpiq

THE DAY PRIOR TO YOUR PROCEDURE. D	ay: Date:
	LIQUIDS ALLOWED: Water, black coffee, apple and white grape
<u> </u>	on or lime Jell-O, chicken or beef broth, and popsicles.
NOT ALLOWED: Solid food, milk, alcohol, or anything RED, ORANGE OR PURPLE.	
STEP 1	Drink one bottle of Clenpiq. Drink Clenpiq right from the bottle.
Crackery Prior MC 53666-67001	Drink 5 cups (8 oz each) of clear liquid. You have from 4:00pm - 10:00pm to complete this step.
CLENPIQ" (sodum picosulfate, magnesium oxide,	Tou have nom 4.00pm - 10.00pm to complete this step.
erd striftwas ciric acid Onal Solution 10 mg/35 g/72 gper 160 mL bottle	
CLENPIQ® is a ready-to-drink oral solution that doesn't need to be diluted.	
CLENDO CL	Repeat Step 1. Stay near a toilet. You have diarrhea, which
STE	
	eventually be a clear yellow liquid.
Drink at least 3 cups (8 oz each) of clear liquid.	

You have until **Midnight** to complete this step.

STEP 4

AT LEAST A WEEK BEFORE YOUR PROCEDURE. Check with your prescribing doctor to make sure it is safe to hold prescription blood thinners. Please get permission to hold Plavix, Coumadin or Warfarin 4 days prior to your procedure. Please get permission to hold Effient, Pradaxa or Xarelto 2 days prior to your procedure.

COLON CLEANSING TIPS:

Walking will help move the solution through yoaur body.

If you feel bloated or your stomach feels "full", drink solution slowly. Stop for 1 hour, then resume.

Continue drinking lots of clear liquids until 4 hours prior to your procedure.

Avoid eating corn 4 days prior to your colon exam.

Anal skin irritation or flare of hemorrhoid inflammation may occur. If this happens, treat it with over-the-counter remedies, such as hydrocortisone cream, baby wipes, Vaseline® or Tucks® – avoid products containing alcohol.

NPO STATUS: During your procedure you will be administered either anesthesia for sedation. For your safety, it is essential you follow these guidelines:

- No GUM, CANDY, MINTS, CIGARETTE SMOKING or TOBACCO USE 4 hours prior to your arrival time.
- DO NOT DRINK ANYTHING 4 HOURS PRIOR TO YOUR COLON / EGD EXAM.

THE DAY OF YOUR PROCEDURE:

- If you take insulin or oral diabetic medication, you are advised to HOLD and DO NOT TAKE on the day
 of your procedure. Bring it with you.
- Take any routine heart, blood pressure or seizure medications with a SMALL sip of water at least two
 hours before your procedure.

YOU MUST HAVE WITH YOU:

- A responsible adult to drive you home. This individual must be present when you check-in and stay at the facility during the procedure or your procedure will be **CANCELLED**.
- A list of all medications you are now taking, including over-the-counter products and herbal supplements.
- A list of any allergies you have.
- If you have a Pacemaker/AICD please bring in your Pacemaker/AICD card.

Midwest Physicians Surgery Center 3601 NE Ralph Powel Road Lee's Summit, MO 64064

Lee's Summit Medical Center 2100 SE Blue Parkway Lee's Summit, MO 64063 Any questions, please contact us at: MIDWEST GI HEALTH - 826-836-2200